

2017年
原著論文

- 1) Masuda S, Shimizu S, Matsuo J, Nishibata Y, Kusunoki Y, Hattanda F, Shida H, Nakazawa D, Tomaru U, Atsumi T, Ishizu A. Measurement of NET formation in vitro and in vivo by flow cytometry. *Cytometry A* 91(8): 822-829, 2017.
- 2) Kawakami T, Okano T, Takeuchi S, Soma Y, Ito F, Ishizu A, Arimura Y, Suzuki K. Elevated moesin mRNA level in skin tissue of patients with polyarteritis nodosa based on real time RT-PCR. *J Dermatol Sci* 87(1): 94-97, 2017.
- 3) Yamada M, Kawakami T, Takashima K, Nishioka Y, Nishibata Y, Masuda S, Yoshida S, Tomaru U, Ishizu A. Establishment of a rat model of thrombosis induced by intravenous injection of anti-phosphatidylserine/prothrombin complex antibody. *Rheumatology (Oxford)* 56(6): 1013-1018, 2017.
- 4) Ishizu A, Tomaru U, Masuda S, Sada KE, Amano K, Harigai M, Kawaguchi Y, Arimura Y, Yamagata K, Ozaki S, Dobashi H, Homma S, Okada Y, Sugiyama H, Usui J, Tsuboi N, Matsuo S, Makino H. Prediction of response to remission induction therapy by gene expression profiling of peripheral blood in Japanese patients with microscopic polyangiitis. *Arthritis Res Ther* 19: 117, 2017.
- 5) Kiuchi S, Tomaru U, Ishizu A, Imagawa M, Kiuchi T, Iwasaki S, Suzuki A, Otsuka N, Deguchi T, Shimizu T, Marukawa K, Matsuno Y, Kasahara M. Expression of cathepsins V and S in thymic epithelial tumors. *Hum Pathol* 60: 66-74, 2017.
- 6) Okano T, Takeuchi S, Soma Y, Suzuki K, Tsukita S, Ishizu A, Suzuki K, Kawakami T. Presence of anti-phosphatidylserine-prothrombin complex antibodies and anti-moesin antibodies in patients with polyarteritis nodosa. *J Dermatol* 44(1): 18-22, 2017.
- 7) Nishioka Y, Yamaguchi M, Kawakami A, Munehiro M, Masuda S, Tomaru U, Ishizu A. Type II natural killer T cells that recognize sterol carrier protein 2 are implicated in vascular inflammation in the rat model of systemic connective tissue diseases. *Am J Pathol* 187(1): 176-186, 2017.

総説・解説、症例報告等

- 1) 石津明洋. 皮膚科領域でみる血管炎の病理組織像からのアプローチ. *日本皮膚アレルギー・接触性皮膚炎学会雑誌* 11(4): 296-299, 2017.